

What is an open breast biopsy?

An open breast biopsy is surgery to remove abnormal breast tissue. It's mostly done when the results of a needle biopsy are uncertain.

To show the location of the abnormal breast tissue, a small wire can be put in the area during a mammogram or ultrasound just before surgery. The wire will guide the doctor to the area to be checked.

The doctor makes a cut in the breast to remove part or all of the abnormal breast tissue. Once the tissue is removed, the doctor will close the cut with stitches.

The breast tissue will be sent to a lab. There it will be examined under a microscope to check for breast cancer. Your doctor may get some answers right away. But it can take up to 1 to 2 weeks to get the final results.

You will be able to go home on the same day as the biopsy. The surgery will leave a scar on your breast that will fade with time. Less often, the surgery may leave a dent in the breast.

How do you prepare for surgery?

Surgery can be stressful. This information will help you understand what you can expect. And it will help you safely prepare for surgery.



Preparing for surgery

- Be sure you have someone to take you home. Anesthesia and pain medicine will make it unsafe for you to drive or get home on your own.
- Tell your doctor ALL the medicines and natural health products you take. Some may increase the risk of problems during your surgery. Your doctor will tell you if you should stop taking any of them before the surgery and how soon to do it.
- If you take aspirin or some other blood thinner, ask your doctor if you should stop taking it before your surgery. Make sure that you understand exactly what your doctor wants you to do. These medicines increase the risk of bleeding.

Make sure your doctor and the hospital have a copy of your goals of care plan. It lets
others know your health care wishes. It's a good thing to have before any type of
surgery or procedure.

Plan Ahead

- **Plan a ride home**. Ask a family member or friend to bring you home after your hospital visit and stay with you for 24 hours..
- **Plan for meals**. Good nutrition is important for your recovery, but cooking may be harder to do right after you get home. Prepare and freeze meals before your surgery or ask friends and family to help you cook or shop for groceries.
- **Get help with chores.** After you get home you may need help. For a short time after your surgery, you may be told to avoid certain physical activities. Plan ahead and arrange to have some help with chores after your surgery.
- **Buy gum or hard candies.** Chewing gum and sucking on hard candies may help get your bowels moving after surgery.

Be as healthy as possible

- **Be active**. Being active before surgery will make it easier for you to be active after surgery. If you're normally active, keep doing your regular activities up to the day of your surgery. If you aren't used to exercising, start slowly. You can start by going for 15 minute walks.
- **Eat healthy**. Eating healthy foods will give your body the nutrients it needs to prepare for and handle the surgery. Eat healthy foods and drink enough fluids in the weeks before your surgery. Your healthcare team will tell you how long before surgery that you need to stop eating and drinking.
- **Manage your medical conditions.** If you have other medical conditions, such as anemia or high or low blood sugar, ask your healthcare team what you need to do to prepare for surgery.
- **Quit tobacco.** You can heal faster and prevent lung problems after surgery if you quit tobacco. Aim to quit smoking and using tobacco and tobacco-like products at least 4 weeks before surgery. Talk to your healthcare team about ways to quit or cut back.
- Don't drink alcohol 24 hours before surgery. Alcohol, cannabis, and other drugs may
 interact with medicines you take before and after surgery. Talk to your healthcare team
 if you need help stopping or cutting back on alcohol, cannabis, or other drugs.

Do the following before your surgery to help you feel better sooner and recover faster.

The day before surgery

- Know the time of your surgery, when you need to be at the hospital, and where you need to go when you get there.
- Follow any instructions the healthcare team or your surgeon gave you for your medicines (including herbal or complementary medicines).

Don't drink alcohol 24 hours before your surgery.



Stop eating at midnight. This is very important!



- Stop eating and drinking all non-clear fluids at midnight. This is very important
- You may DRINK CLEAR FLUIDS ONLY up to 3 hours before surgery. This includes BLACK coffee, tea, Gatorade, apple juice and water.

Morning of surgery

Plan to arrive at the hospital on time.

- Follow any instructions provided
- Bring all of the things you packed for your hospital stay **if** you are being admitted.
- Follow **all** other instructions the healthcare team gave you to prepare for surgery.
- Bring your CPAP if you use one



•

What happens on the day of surgery?

If your doctor told you to take your medicines on the day of surgery, take them with only a sip of water.

- Take a bath or shower before you come in for your surgery. Do not apply lotions, perfumes, deodorants, or nail polish.
- Do not shave the surgical site yourself.
- Take off all jewellery and piercings. And take out contact lenses, if you wear them.
- Bring a comfortable, supportive bra with you.
- At the hospital



Bring a picture ID health card

- The area for surgery is often marked to make sure there are no errors. If needed, you will get an ultrasound to mark the area.
- You will be kept comfortable and safe by your anesthesia provider.

- When should you call your doctor?
- You have questions or concerns.
- You don't understand how to prepare for your surgery.
- You become ill before the surgery (such as fever, flu, or a cold).
- You need to reschedule or have changed your mind about having the surgery.

Open Breast Biopsy: What to Expect at Home

Your Recovery

An open breast biopsy is surgery to remove abnormal breast tissue. The breast tissue will be sent to a lab, where a doctor will look at the tissue under a microscope to check for breast cancer.

Your doctor may have some answers right away. But it can take up to 1 to 2 weeks to get the final results. Your doctor will discuss the results with you.

For a few days after the surgery, you will probably feel tired and have some pain. The skin around the cut (incision) may feel firm, swollen, and tender. The area may be bruised. Tenderness should go away in about a week, and the bruising will fade within two weeks. Firmness and swelling may last 6 to 8 weeks.

Your incision may have been closed with strips of tape or stitches. If you have strips of tape on the incision, leave the tape on for a week or until it falls off. If you have stitches they will dissolve in 3-4 weeks

This care sheet gives you a general idea about how long it will take for you to recover. But each person recovers at a different pace. Follow the steps below to get better as quickly as possible.

How can you care for yourself at home?



Activity

- Rest when you feel tired. Getting enough sleep will help you recover.
- Try to walk each day. Start by walking a little more than you did the day before. Bit by bit, increase the amount you walk. Walking boosts blood flow and helps prevent pneumonia and constipation.

- For 2 weeks, avoid strenuous activities that put pressure on your chest or that involve vigorous movement of your upper body and arm on the side of the biopsy. Examples of these might include strenuous housework, holding an active child, jogging, or aerobic exercise.
- For 2 weeks, avoid lifting anything that would make you strain. This may include heavy grocery bags and milk containers, a heavy briefcase or backpack, cat litter or dog food bags, a vacuum cleaner, or a child.
- Ask your doctor when you can drive again.



Diet

• Start off with light foods such as soup, toast and tea. Avoid dairy and fatty foods as it may cause stomach upset



Medicines

- Your doctor will tell you if and when you can restart your medicines. He or she will also give you instructions about taking any new medicines.
- If you take aspirin or some other blood thinner, ask your doctor if and when to start taking it again. Make sure that you understand exactly what your doctor wants you to do.
- If the doctor gave you a prescription medicine for pain, take it as prescribed.
- If you are not taking a prescription pain medicine, ask your doctor if you can take an over-the-counter medicine.
- If your doctor prescribed antibiotics, take them as directed. Do not stop taking them just because you feel better. You need to take the full course of antibiotics.



Incision care

- If you have strips of tape on the incision, leave the tape on for a week or until it falls off.
- Wash the area daily with warm, soapy water, and pat it dry. Don't use hydrogen peroxide or alcohol, which can slow healing. You may cover the area with a gauze bandage if it weeps or rubs against clothing. Change the bandage every day.
- Keep the area clean and dry.

Other instructions

• For the first 14 days after surgery, wear a supportive bra all the time, even at night.

Discharge

How you may feel after your operation

- You may feel nauseated, light headed dizzy or sleepy for up to 24 hours after your surgery
- You may have a slight temperature for 2 days
- You may feel tired for a few days after your surgery

For 24 hours after your surgery do not.... Drive a vehicle

Operate heavy equipment Drink alcohol or smoke

Make any important or legal decisions

At Home

Slowly get more active.

- Eat healthy and drink fluids as you did before surgery (unless your healthcare team gives you other instructions).
- Drink oral nutritional supplements (like Ensure Protein Max) for the first month after surgery to give you extra protein and help you heal faster.
- Take less pain medicine as your pain gets better.
- Take your blood thinner and other medicines as you were told.

What is anesthesia?

Anesthesia controls pain. And it keeps all your organs working normally during surgery or another kind of procedure.

Anesthesia can relax you. It can also make you sleepy or forgetful. Or it may make you unconscious. It depends on what kind you get.

Your anesthesia provider (anesthesiologist) will make sure you are comfortable and safe during the procedure or surgery.

There are different types of anesthesia.

- **Local anesthesia.** This type numbs a small part of the body. Doctors use it for simple procedures.
- You get a shot in the area the doctor will work on.

- o You will feel some pressure during the procedure.
- o You may stay awake. Or you may get medicine to help you relax or sleep.
- **Regional anesthesia.** This type blocks pain to a larger area of the body. It can also help relieve pain right after surgery. And it may reduce your need for other pain medicine after surgery. There are different types. They include:
- Peripheral nerve block. This is a shot near a specific nerve or group of nerves. It blocks pain in the part of the body supplied by the nerve. This is often used for procedures on the hands, arms, feet, legs, or face.
- Epidural and spinal anesthesia. This is a shot near the spinal cord and the nerves around
 it. It blocks pain from an entire area of the body, such as the belly, hips, or legs.
- **General anesthesia.** This type affects the brain and the whole body. You may get it through a small tube placed in a vein (IV). You are unconscious and will not feel pain. During the surgery, you will be comfortable. Later, you will not remember much about the surgery.

What type will you have?

The type of anesthesia you have depends on many things, such as:

- The type of surgery or procedure and the reason you are having it.
- Test results, such as blood tests.
- How worried you feel about the surgery.
- Your health. Your doctor and nurses will ask you about any past surgeries. They will ask about any health problems you may have, such as diabetes, lung or heart disease, or a history of stroke. They will want to know if you take medicine, such as blood thinners. Your doctor may also ask if any family members have had any problems with anesthesia. You will talk with your anesthesia provider about your options. In many cases, you may be able to choose the type of anesthesia you have.

What can you expect after having anesthesia?

Right after the surgery, you will be in the recovery room. Nurses will make sure you are comfortable. As the anesthesia wears off, you may feel some pain and discomfort from your surgery.

Tell someone if you have pain. Pain medicine works better if you take it before the pain gets bad.

You may feel some of the effects of anesthesia for a while. It takes time for the effects of the medicine to completely wear off.

- If you had local or regional anesthesia you may feel numb and have less feeling in part
 of your body. It may also take a few hours for you to be able to move and control your
 muscles as usual.
- When you first wake up from general anesthesia, you may be confused. Or it may be hard to think clearly. This is normal.
- Don't do anything for 24 hours that requires attention to detail. This includes going to work, making important decisions, or signing any legal documents.
 Other common side effects of anesthesia include:
- Nausea and vomiting. This does not usually last long. It can be treated with medicine.
- A slight drop in body temperature. You may feel cold and shiver when you first wake up.
- A sore throat, if you had general anesthesia.
- Muscle aches or weakness.
- Feeling tired.

You may go home the same day of surgery or you may stay in the hospital. Your doctor will check on your recovery from the anesthesia. He or she will answer any questions you may have.

For answers to frequently asked questions about your anesthetic, please visit https://www.cas.ca/en/about-cas/advocacy/anesthesia-faq

When should you call for help?

Call 911 anytime you think you may need emergency care. For example, call if:

- You passed out (lost consciousness).
- You have chest pain, are short of breath

Call your doctor or seek immediate medical care if:

- You have pain that does not get better after you take pain medicine.
- You have signs of infection, such as:
- Increased pain, swelling, warmth, or redness.
- Red streaks leading from the incision.
- $^{\circ}$ Pus draining from the incision.
- A fever.
- Bright red blood has soaked through the bandage over your incision.